



Delaware Military Academy

2019 National Blue Ribbon School
20 years **STRONG** in developing Citizens and Scholars
to lead our Nation through the 21st century!



Self-Administration of Asthma Inhaler Student Agreement

Name: _____

Grade: _____

Inhaled Medication: _____

Date: _____

I agree to:

- Follow my prescribing health professional's medication order.
- Use correct medication administration technique.
- Not allow anyone else to use my medication under any circumstances.
- Keep the medication with me in school and on field trips.
- Inform the school nurse of the time and reason for taking the inhaler.
- Notify (or have someone else notify) the school nurse immediately if the following occurs:
 - My symptoms continue to get worse after taking the medication.
 - My symptoms reoccur within 2-3 hours after taking the medication.
 - I think I might be experiencing side effects from my medication.
 - Other _____
- I understand that permission for self-administration of medication may be discontinued if am unable to follow the safeguards established above.

Signature of Student

Date

Signature of Parent/Guardian/Relative Caregiver

Date

-
- Student verbalizes dose _____
 - Student demonstrates proper technique
 - Removes cap and shake if applicable
 - Attaches spacer if applicable
 - Breathes out slowly
 - Presses down inhaler to release medication
 - Breathes in slowly
 - Holds breath for 10 seconds
 - Repeats as directed
 - Student verbalizes safe use
 - Student verbalizes symptoms/signs of when medication is needed & when to notify school nurse
 - Parent permission to self-administer

The student has demonstrated knowledge about the proper use of his/her medication and necessary permissions (parent and licensed healthcare provider) are on file.

Signature of School Nurse

Date

CITIZENSHIP • LEADERSHIP • SCHOLARSHIP

112 Middleboro Road • Wilmington, DE 19804 • 302-998-0745 • 302-998-3521 Fax • www.demilacad.org

THE DELAWARE MILITARY ACADEMY IS AN EQUAL OPPORTUNITY EMPLOYER. IT DOES NOT DISCRIMINATE ON THE BASIS OF RACE, COLOR, RELIGION, NATIONAL ORIGIN, SEX, SEXUAL ORIENTATION, GENDER IDENTITY, MARITAL STATUS, DISABILITY, AGE, GENETIC INFORMATION, HOUSING/HOMELESS STATUS OR VETERAN STATUS IN EMPLOYMENT, OR ITS PROGRAMS AND ACTIVITIES.



Asthma Action Plan

General Information:

DOB ____ / ____ / ____

- Name _____
- Emergency contact _____ Phone numbers _____
- Physician/Health Care Provider _____ Phone numbers _____
- Physician Signature _____ Date _____

Severity Classification	Triggers	Exercise
<input type="radio"/> Mild Intermittent <input type="radio"/> Moderate Persistent <input type="radio"/> Mild Persistent <input type="radio"/> Severe Persistent	<input type="radio"/> Colds <input type="radio"/> Smoke <input type="radio"/> Weather <input type="radio"/> Exercise <input type="radio"/> Dust <input type="radio"/> Air pollution <input type="radio"/> Animals <input type="radio"/> Food <input type="radio"/> Other	1. Pre-medication (how much and when) _____ _____ 2. Exercise modifications _____

Green Zone: Doing Well

Peak Flow Meter Personal Best = _____

Symptoms

- Breathing is good
- No cough or wheeze
- Can work and play
- Sleeps all night

Peak Flow Meter

More than 80% of personal best or _____

Control Medications

Medicine	How Much to Take	When To Take It
_____	_____	_____
_____	_____	_____

* Both the Healthcare Provider and the Parent/Guardian see that the child has demonstrated the skills to carry and self-administer their quick relief inhaler, including when to tell an adult if symptoms do not improve after taking the medicine.

Yellow Zone: Getting Worse

Contact Physician if using quick relief more than 2 times per week.

Symptoms

- Some problems breathing
- Cough, wheeze or chest tight
- Problems working or playing
- Wake at night

Peak Flow Meter

Between 50 to 80% of personal best or _____ to _____

Continue control medicines and add:

Medicine	How Much to Take	When To Take it
_____	_____	_____
_____	_____	_____

IF your symptoms (and peak flow, if used) return to Green Zone after one hour of the quick relief treatment, THEN

- Take quick-relief medication every 4 hours for 1 to 2 days
- Change your long-term control medicines by _____
- Contact your physician for follow-up care

IF your symptoms (and peak flow, if used) DO NOT return to the GREEN ZONE after 1 hour of the quick relief treatment, THEN

- Take quick-relief treatment again
- Change your long-term control medicines by _____
- Call your physician/Health Care Provider within _____ hours of modifying your medication routine

Red Zone: Medical Alert

Ambulance/Emergency Phone Number: _____

Symptoms

- Lots of problems breathing
- Cannot work or play
- Getting worse instead of better

Peak Flow Meter

Between 0 to 50% of personal best or _____ to _____

Continue control medicines and add:

Medicine	How Much to Take	When To Take It
_____	_____	_____
_____	_____	_____

Go to the hospital or call for an ambulance if

- Still in the red zone after 15 minutes
- If you have not been able to reach your physician/health care provider for help
- _____

Call an ambulance immediately if the following danger signs are present

- Trouble walking/talking due to shortness of breath
- Lips or fingernails are blue